

Evaluation of Factors Affecting Mortality in Patients with Severe Maxillofacial Injuries in Sulaimaniyah City/KRG

Farman Salih Jamal^{1*}, Bayad Jaza Mahmood²

^{1*}BDS., KHCMS Candidate Maxillofacial Surgery, Sulaimani Center, Kurdistan Region-Iraq.

²BDS., FKHCMS., AACME., Assistant. Prof., Head of Maxillofacial Surgery, College of Dentistry, University of Sulaimaniyah, Kurdistan Region-Iraq.

Corresponding author:

BDS., KHCMS Candidate Maxillofacial Surgery, Sulaimani Center, Kurdistan Region-Iraq

Cite this paper as: Farman Salih Jamal, Bayad Jaza Mahmood.(2024) Evaluation of Factors Affecting Mortality in Patients with Severe Maxillofacial Injuries in Sulaimaniyah City/KRG *Frontiers in Health Informatics*, 13(8) 6572-6578

ABSTRACT

Background and objectives: Traumatic injuries continue to be a major contributor to mortality, and long-term disability all over the world. Facial injuries impact one-third of severely injured patients. Occasionally, standalone maxillofacial trauma is linked with fatality, however in patients with concurrent injuries, the death rate increases. This research focused on analyzing and evaluating facial injuries and factors affecting mortality among patients admitted to the emergency unit due to traumatic injuries.

Methods: A Prospective Cross-sectional methodology was used in this research; consisting of a total of 105 patients of both sexes including 81 males and 24 females with maxillofacial injuries were observed until carrying out the hospital or dying upon arrival or during their hospital stay between January 2024 and January 2025 at the emergency department of Shar Teaching Hospital in Sulaimaniyah city, Kurdistan region, Iraq. Measurements of Injury Severity Score and Glasgow Coma Scale were statistically analyzed to determine the factors associated with mortality. Injury Severity Score is evaluated by the process of converting injury codes in the International Classification System of Diseases, tenth edition (ICD-10).

Result: During a one-year observation period, 105 patients were evaluated. The average age (mean) of individuals with maxillofacial injuries was 28.1 ± 14.9 years, and 77% of them were male. Road traffic accidents including (car accidents and motorcycle collisions) were the most prevalent etiological cause of facial injuries regardless of gender. The research findings reveal a significant correlation between mortality and AIS-ISS scores and GCS.

keywords: Maxillofacial injury, Mortality, Glasgow Coma Scale, Injury Severity Score

Introduction

Traumatic injuries continue to be a major contributor to mortality, and long-term disability all over the world. Facial injuries impact one-third of severely injured patients [1,2]. Traumatic injuries have been established as one of the significant population health problems in developed nations in addition to nations with low yearly income [3]. A considerable proportion of trauma patients (40-60%) pass away caused by injuries before arriving at the hospital [4]. The incidence of traumatic injuries is more widespread among young individuals, predominantly men (70%-80%) which leads to a substantial economic burden for society because life expectancy is reduced as a result of fatality and disability [5]. Given that the facial bones are anatomically positioned in distinct balance and their close and strong relation to the brain, traumatic facial traumas are of critical significance [6]. Globally, the predominant factors of maxillofacial trauma are vehicle collisions, then falls, assault, and other causes [7].

Studies in the existing literature have established that maxillofacial injuries is mainly found in men within the age range of 21 to 30. The ratio of male to female has been established between 2:1 and 8:1 in the research literature[7,8]. Facial injuries are typically responsible for nonreversible consequences or fatality in individuals, particularly while they are correlated with other concurrent injuries. Head injuries and thoracic injuries are usually linked with maxillofacial bone fractures in patients with extensive traumatic injuries [9]. Head injury, chest injury, and Abbreviated Injury Scale (AIS), Injury Severity Score (ISS) are recognized as stable and unconventional markers in increasing the rates of mortality among patients with polytraumatic injuries[10]. According to existing research, isolated facial injuries are occasionally associated with death. At the same time patients with concomitant traumatic injuries, the rates of mortality are significantly elevated [11]. Based on the published studies, the prevalence rate of head injuries in patients with maxillofacial injuries was up to 20–70% contingent upon the socioeconomic and environmental situation [12]. Most often, head injuries correspond with severe maxillofacial injuries such as midface fractures and/or naso-orbito-ethmoid (NOE) fractures frontal bone fractures, or sinus entanglement; head injuries are less uncommonly correlated with isolated fractures of the mandible.

Besides the head participating, other concomitant injuries can be incorporated such as chest, abdomen, and extremities [13]. The scores of the Abbreviated Injury Scale (AIS), which is anatomically derived, were suggested in 1971 and have continued brought up numerous times. In 1974, Baker et al. established the Injury Severity Score (ISS) derived from the supposition that the degree of intensity of the injuries and probability of death could be exhibited by the cumulative sum of the squaring of the triple highest AIS points for three separate body parts. The AIS-ISS approach has been established as the most broadly adopted survival indicator guide in medical outcomes [14,15].

Materials and Methods

A Prospective Cross-sectional methodology was used in this research; consisting of a total of 105 patients of both sexes including 81 males and 24 females with maxillofacial injuries were observed until carried out to the hospital or dying on arrival or during their hospital stay between January 2024 and January 2025 at the emergency department of Shar Teaching Hospital in Sulaimanyah city, Kurdistan region, Iraq. The scientific study protocol is authorized by the Ethical Committee of the Kurdistan Higher Council of Medical Specialties (KHCMS) Ethical Committees.

Injuries to the Maxillofacial region involve damage to the soft and/or hard tissue of the face and jaws. Each patient record is evaluated individually. The following parameters were gathered and analyzed: patient demography, etiology and mechanism of trauma, anatomical body region and intensity of the injury, concomitant injuries, and clinical outcome. Maxillofacial fractures and head injuries were evaluated by X-ray and/or computed tomography (CT) scan of the face and head. CT/X-ray investigations were carried on further to the thorax, abdomen, spine, and extremities when multiple injuries were recognized.

The severity and intensity of the injury of regional anatomy were analyzed by the following commonly used grading methods: (AIS) The Abbreviated Injury Severity Score, (ISS) The Injury Severity Score, and (GCS) Glasgow Coma Score. AIS is a global scoring evaluation system that categorizes any injury by anatomical region on the 6-point ordinal scale for grading injuries 1=minor, 2=moderate, 3=serious, 4=severe, 5=critical, 6=maximal. The Values of the Injury Severity Score (ISS) are evaluated by the process of converting injury codes in the International Classification System of Diseases, tenth edition (ICD-10). The injury severity score (ISS) minor=1-8, moderate=9-15, severe=16-24, critical=25-75 was established according to the records. The AIS-ISS score was evaluated for any individual dependent on diagnoses.

Patients with concomitant injuries were graded by the cumulative sum of the squares of the three greatest AIS grades for three separate body parts. Information regarding the state of consciousness and awareness based on the GCS was obtained from the data notes given by the first neurosurgeon who evaluated the patient's neurological states. The difference between groups concerning parameters such as AIS, ISS, and GCS were estimated. Further analysis was conducted to recognize the variables that were significantly correlated with death rates. The data were gathered and arranged via Excel spreadsheet software (Microsoft). Before the numerical data analysis, the data were checked to ensure the required data quality.

Statistical Analysis

The recorded data was transferred into the GraphPad Prism 9.0 Statistical program. The numerical data process was conducted by GraphPad Prism 9.0.2 program and was used for the visualization of the results. Quantitative continuous variables were revealed as mean, and standard deviation. Median range categorical ordinal variables and categorical nominal data were introduced in the form of percentages (%). The variation between categorical data was evaluated with the unpaired t-test, the Mann-Whitney U test. Several logistic regression evaluation was used to analyze the correlation between variables including (gender, age etiological factor of injury AIS-ISS), and death rate in individuals with multiple traumas, and the Odds ratio (OR) was determined with a 95% (CI) confidence interval for the variables. A p-value of <0.05 was regarded to be statistically significant.

Results

During the 12-month observation period, 105 patients were collected and evaluated. The patient's mean age included in this study was (28.2, SD \pm 14.9 years), ranging from 3 to 80 years. There were 81 males (77%) and 24 females (23%) with a gender proportion of 3,3:1. The majority of trauma occurred amongst the 21-30 year (young) age group. Road traffic accidents including (car accidents, motorcycle accidents, and pedestrian accidents) were the most prevalent etiological cause (74%) with statistical significance in comparison to the other groups ($p < 0.001$). The second most common cause of injuries was fall from height (11%). The other factors (15%) include assault, work-related injuries, blast injuries, sport-related injuries, and unknown causes. The evaluation of data revealed that there was no relevant statistical variation in the population among patients who died and those who survived involving gender, age, etiological factors of injuries (Table 1 and Figure 1).

Table 1. The demographic of maxillofacial injured patients, gender, and mechanism of the trauma of patients those who survived, and those who nonsurvived.

Parameters Maxillofacial Injured patients The rate in percentage (%)

Parameters		Maxillofacial Injured patients	The rate in percentage (%)
Age (yr) (Mean \pm SD)		28.1 \pm 14.9	
Gender	Male	81	77.2
	Female	24	22.8
Mechanism of injuries	Road traffic accident	78	74.3
	Falling accident	11	10.5
	Others	16	15.3
Death		5	4.8




Figure 1. The chart illustrates the age demographics range of patients

In this study, Among all patients only (32.7%)patients suffered isolatedmaxillofacial injuries, while (67.3%)patients had concomitant injuries in another body region. In polytraumatic patients, The most frequently injured body anatomical regions associatedwith facial fractures were the head injuries (49.6%), followed by the chest injuries (27.7%), then the extremities(17.2%). The face-AIS score corresponded to minor in 59 % of patients, moderate in 21 %, serious in 12%, severe in 5%, critical 2%, and maximal in the remaining 1%. The median range (AIS-ISS) was 5 (1-75) about 80 %of patients had an ISS between 1 and 8. The mean GCS measurement of the patients upon admission to the emergency unit was observed to be 12.7 ± 2.9 . A GCS score ≤ 9 was recognized in (12.4%) of the individuals. The lowest possible score was 3.Multiple predictor regression modeling reveals that facial-AIS(Odd ratio: 2.14(95% CL: 1.119—3.166)) and GCS (Odd ratio: 12.17(95% CL: 12.13—13.26)). The overall mortality rate for the patients in this study was (4.8%) although there was a male predominance in the patients with maxillofacial injuries, there was no gender-based variationregarding survival.Table 2 shows thedetails of differentiating the intensity indicators among the survived and nonsurvived patients and reveals that patients who died had considerably lower GCS and substantially higher ISS.

Table 2. Two sample-based data analyses (median values) of comparing and evaluating intensity indicators of facial injured patients who survived with those who did not survived.

Variable		Total (N=105)		Survivors (N=100)		Non Survivors (N=5)	
		Median	Range	Median	Range	Median	Range
Location of injury	Face AIS	1	(1-6)	1	(1-3)	4	(2-6)
	Head AIS	2	(1-5)	2	(1-5)	4	(2-5)
	Chest AIS	1	(1-4)	1	(1-4)	2	(1-4)
Injury Severity Score (ISS)		5	(1-75)	5	(1-75)	19.5	(5-75)
Glasgow Coma Scale (GCS)		14	(3-15)	14	(3-15)	4	(3-15)

Discussion

Traumatic injuries rank as one of the key public health concerns across the world and are a substantial contributor to death and disability, particularly among the young demographic[16]. Maxillofacial trauma results in serious healthcare, economic, and societal outcomes that are influenced by considerable appearance and efficiency disabilities, as well as psychological disturbances, social impairments, and high financial burden associated with medical care and recovery. Maxillofacial injuries differ in form, intensity, and signs and symptoms. They can also be related to concurrent injuries in the other body regions. The occurrence of numerous critical injuries in the facial region is correlated with a greater risk of complications, clinical outcomes, and even fatality[17]. Polytrauma is typically more critical, and the possibility of death is higher in the case of head injuries, with 70.77% of mortality caused by polytrauma who suffered head injuries [18,19].

The result of the present study is similar to other previous studies revealing that facial injuries occurred in age related curve with the highest incidence seen between ages 21 to 30 years. In addition, children under 11 are included in 4% to 10% of all facial injuries [20]. At the same time in this research the ratio of males to females significantly higher were males 3 times more than females 77% and 23% of male and female that in sequence, these findings were nearly compatible with most of the research on this topic, This is characteristic to the reason that male furthermore engaged in external occupation moreover contact to assault and aggression interaction. In addition, the a higher proportion of male car drivers or motorcycle drivers than female. This ratio seems to be less prominent in developed nations due to greater involved outdoor activity of female [21,22]. In our patient groups, the highly frequent etiologic factor of facial injuries was traffic collision 74% as reported in other studies from other countries of the world[23-25].

In addition, in the current research, the next most frequent cause of injuries was falls from height 11% which was attributed to inadequate safety protocols in the workplace. The other factors in 15% include assault, blast injuries, sport-related injuries, and unknown causes[26]. Concomitant injuries are common in maxillofacial injured patients, they occurred in 67.3% of all clinical cases, there is a significant interrelation between the occurrence of accompanying injuries and the mechanism of traumatization[27]. In all instances, the primary cause of death was correlated with concomitant injuries and not directly caused by facial injuries, head injuries, and thoracic injuries are prevalent in the current study where head injuries occur in (49.6%), followed by chest injuries (27.7%), then other concomitant injuries [28-30]. Consequently, the close positioning of the head and chest correlatively to the facial region enhances their chance of injury. This basic clinical observation is confirmed by the evidence of our study.

The overall mortality rate for the patients in this study was 4.8% which is near to other studies[27,29]. Data analysis revealed that high AIS-ISS, reduced GCS, and associated chest injuries were considerably high in individuals who died identical to previous studies[27,30]. We observed a significant connection between the maxillofacial injuries severity and head injuries. Furthermore, we identified that a combination of multiple facial bone fractures correlated with more critical head injuries, as determined by the GCS and AIS-ISS. Interestingly the current study and other literature found a strong and linear correlation between AIS-ISS and survival rate, but generally, maxillofacial trauma presents a lower AIS-ISS score than head and neck injuries[17,31].

Conclusion

In conclusion, facial traumatic injuries are a frequent form of which leads to moderate to serious injuries. The majority of maxillofacial injured patients are young males that injured by road traffic accidents or falls from height. Furthermore, the rate of mortality among patients with facial injuries was 4.8% overall. In addition, the most important factors affecting survival rate are associated with injuries to another part of the body such as head injuries, chest injuries, and others that are determined by GCS, and AIS-ISS scores. Moreover, the rate of mortality is greater in patients with severe facial injuries such as maxillary lefort III fracture, and pan-facial fracture than in those with mandibular fracture.

Interest conflict

All authors have no conflict of interest related to the release or publication of article.

Consent for publications

The author checked and confirmed the final version of the manuscript for publication.

Ethics approval and consent to participate

The Research protocol approved by the Ethical Committee of the Kurdistan Higher Council of Medical Specialities (KHCMS) Ethical Committees

REFERENCES

1. Gonsaga RAT, Brugugnolli ID, Fraga GP. Comparison between two mobile pre-hospital care services for trauma patients. *World Journal of Emergency Surgery*. 2012;7(1):1.
2. EsmerE, DelankKS, SiekmannH, SchulzM, DerstP. Facial injuries in polytrauma—which injuries can be expected? A retrospective evaluation from the TraumaRegister DGU(R) Notfall Rettungsmed. 2016; 19:92–98
3. Zandi M, SeyedHoseini SR (2013) The relationship between head injury and facial trauma: a case-control study. *Oral Maxillofac Surg* 17(3):201–207
4. Smith GS, Barss P. Unintentional injuries in developing countries: the epidemiology of a neglected problem. *Epidemiologic reviews*. 1991;13(1):228–66.
5. Akbari GA, Mohammadian A. Comparison of the RTS and ISS scores on prediction of survival chances in multiple trauma patients. *Injury*. 2012;6:11.
6. Maas AI, Stocchetti N, Bullock R. Moderate and severe traumatic brain injury in adults. *The Lancet Neurology*. 2008;7(8):728–41.
7. Schaftenaar E, Bastiaens GJH, Simon ENM, Merckx MAW. Presentation and management of maxillofacial trauma in Dar Es Salaam, Tanzania. *East Afr Med J*. 2009;86(6):254–8
8. Farias IPSE, Bernardino ÍM, Nóbrega LMD, Gempel RG, D'Avila S. Maxillofacial trauma, etiology and profile of patients: an exploratory study. *Acta Ortop Bras*. 2017 Nov-Dec;25(6):258–261
9. Almasri M. Severity and causality of maxillofacial trauma in the Southern region of Saudi Arabia. *Saudi Dent J*. 2013;25(3):107–10.
10. Prathigudupu RS, Tiwari RVC, Mathew P, Roy B, Sadique S, Tiwari H (2018) Mortality in maxillofacial trauma – a review. *Saudi J Oral Dent Res* 3(7):219–223
11. Paydar S, Moghaninasab A, Asiaei E, Jahromi GSF, Bolandparvaz S, Abbasi H. Outcome of patients underwent emergency department thoracotomy and its predictive factors. *Emergency*. 2014;2(3):125.
12. Arajärvi E, Lindqvist C, Santavirta S, Tolonen J, Kiviluoto O (1986) Maxillofacial trauma in fatally injured victims of motor vehicle accidents. *Br J Oral Maxillofac Surg* 24(4):251–257.
13. You N, Choi MS, Roh TH, Jeong D, Kim SH (2018) Severe facial fracture is related to severe traumatic brain injury. *World Neurosurg* 111:e47–e52.
14. Haug RH, Wible RT, Likavec MJ, Conforti PJ (1991) Cervical spine fractures and maxillofacial trauma. *J Oral Maxillofac Surg* 49(7):725–729.
15. Baker SP, O'Neill B, Haddon W Jr, Long WB (1974) The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 14(3):187–196.
16. Chen C, Zhang Y, An JG, He Y, Gong X (2014) Comparative study of four maxillofacial trauma scoring systems and expert score. *J Oral Maxillofac Surg* 72(11):2212–2220
17. Segui-Gomez M, MacKenzie EJ. Measuring the public health impact of injuries. *Epidemiologic Reviews*. 2003;25(1):3–19.
18. Bagheri SC, Dierks EJ, Kademani D, Holmgren E, Bell RB, Hommer L, Potter BE (2006) Application of a facial injury severity scale in craniomaxillofacial trauma. *J Oral Maxillofac Surg* 64(3):408–414.
19. Lavoie A, Moore L, LeSage N, Liberman M, Sampalis JS (2004) The new injury severity score: a more accurate predictor of in-hospital mortality than the Injury Severity Score. *J Trauma* 56(6):1312–1320.
20. Domingues Cde A, de Sousa RM, Nogueira Lde S, Poggetti RS, Fontes B, Muñoz D (2011) The role of the New Trauma and Injury Severity Score (NTRISS) for survival prediction. *Rev Esc Enferm USP* 45(6):1353–1358.
21. Moore EE, Feliciano DV, Mattox KL. *Trauma*. 5th ed. McGraw-Hill Professional: New York;2003. p.149–165.
22. Oji C. Jaw fractures in Enugu, Nigeria, 1985–95. *Br J Oral Maxillofac Surg* 1999; 37: 106–109.
23. Meyer U, Benthous S, Du Chesne A, Wannhof H, Zöllner B, Joos U. Examining patients with facial skull fractures

- from an etiological and legal perspective. *Mund KieferGesichtschir* 1999; 3: 152-157.
23. Bataineh AB. Etiology and incidence of maxillofacial fractures in the north of Jordan. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1998; 86: 31-35.
 24. Bailey BJ, Johnson JT, Newlands SD, et al. *Head & Neck Surgery: Otolaryngology*. 4th ed., Hagerstown, MD: Lippincott Williams & Wilkins; 2005. p. 935-936.
 25. Oikarinen K, Schutz P, Thalib L, Sándor GK, Clokie C, Meisami T, Safar S, et al. Differences in the etiology of mandibular fractures in Kuwait, Canada, and Finland. *Dent Traumatol* 2004; 20: 241-245.
 26. Barss P, Addley K, Grivna M, Stanculescu C, Abu-Zidan F. Occupational injury in the United Arab Emirates: epidemiology and prevention. *Occup Med (Lond)*. 2009; 59:493-498.
 27. Shumynskyi I, Gurianov V, Kaniura O, Kopchak A. Prediction of mortality in severely injured patients with facial bone fractures. *Oral Maxillofac Surg*. 2021.
 28. Eid HO, Abu-Zidan FM. Biomechanics of road traffic collision injuries: a clinician's perspective. *Singapore J*. 2007; 48:693-700
 29. Işık D, Go"nu" llu" H, Karadaş S, Koc,ak OF, Keskin S, Garca MF, et al. Presence of accompanying head injury in patients with maxillofacial trauma. *Ulus Travma AcilCerrahi Derg*. 2012; 18:200-206.
 30. Rajandram RK, Syed Omar SN, Rashdi MF, Abdul Jabar MN. Maxillofacial injuries and traumatic brain injury—a pilot study. *Dent Traumatol*. 2014; 30:128-132. Rajandram RK, Syed Omar SN, Rashdi MF, Abdul Jabar MN. Maxillofacial injuries and traumatic brain injury—a pilot study. *Dent Traumatol*. 2014; 30:128-132.
 31. Down KE, Boot DA, Gorman DF. Maxillofacial and associated injuries in severely traumatized patients: implications of a regional survey. *Int J Oral Maxillofac Surg*. 1995;24(6):409-12.